

AD-A145 357

SPECTROPHOTOMETRIC STUDIES OF IODINE COMPLEXES IN AN  
ALUMINUM CHLORIDE ... (U) STATE UNIV OF NEW YORK AT  
BUFFALO DEPT OF CHEMISTRY Z J KARPINSKI ET AL. JUL 84  
SUNYBUF/DC/TR-20 N00014-79-C-0682

1/1

UNCLASSIFIED

F/G 20/6

NL


END

DATE

FORMED

16 84

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS - 1963 - A

4

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SUNYBUF/DC/TR-20	2. GOVT ACCESSION NO. AD-A145 357	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Spectrophotometric Studies of Iodine Complexes In An Aluminum Chloride - Butylpyridinium Chloride Ionic Liquid		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Zenon J. Karpinski and Robert A. Osteryoung		8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0682
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry State University of New York at Buffalo		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-715
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Arlington, VA 22217		12. REPORT DATE July, 1984
		13. NUMBER OF PAGES 23
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release: Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for Publication in Inorganic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Iodine Complexes; Spectrophotometry; Molten Salts		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Iodine, iodine chloride, tetraethylammonium iodide and tetraethylammonium triiodide solutions in aluminum chloride - N-(1-butyl)pyridinium chloride ambient temperature ionic liquids have been studied by u.v. - visible spectrophotometry. The different oxidation states of iodine show a distinct dependence on the solvent acidity. Iodine(+1) exists in the form of $ICl$ in acidic melts and in the form of $ICl_2^-$ in basic $AlCl_3$ -BuPyCl mixtures. Molecular iodine does not form strong complexes in acidic or neutral solutions. Triiodide ion is stable		

DTIC  
SEP 07 1984  
E

84 - 09 06 063

AD-A145 357

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

>in neutral butylpyridinium tetrachloroaluminate, but not in acidic melts. In solutions of iodine or triiodide in basic  $\text{AlCl}_3$ -BuPyCl ionic liquids equilibrium mixtures of  $\text{I}_2\text{Cl}^-$ ,  $\text{I}_3^-$ ,  $\text{ICl}_2^-$  and  $\text{I}^-$  are formed.

The charge transfer band observed for iodide solutions in neutral and basic melts reveal the formation of iodide ion - butylpyridinium cation ion pairs. Similar bands in pure basic  $\text{AlCl}_3$ -BuPyCl mixtures indicate the association of chloride ion and butylpyridinium<sup>3</sup> cation.

Accession For	
NTIS GSA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0682

Task No. NR-051-715

TECHNICAL REPORT NO.20

Spectrophotometric Studies of Iodine Complexes  
In An Aluminum Chloride - Butylpyridinium Chloride  
Ionic Liquid

by

Zenon J. Karpinski and Robert A. Osteryoung

Prepared for Publication

in

Inorganic Chemistry

Department of Chemistry  
State University of New York at Buffalo  
Buffalo, New York 14214

July, 1984

Reproduction in whole or in part is permitted for any purpose of the  
United States Government

Approved for Public Release: Distribution Unlimited

ABSTRACT

Iodine, iodine chloride, tetraethylammonium iodide and tetraethylammonium triiodide solutions in aluminum chloride - N-(1-butyl)pyridinium chloride ambient temperature ionic liquids have been studied by u.v. - visible spectrophotometry. The different oxidation states of iodine show a distinct dependence on the solvent acidity. Iodine(+1) exists in the form of  $\text{ICl}$  in acidic melts and in the form of  $\text{ICl}_2^-$  in basic  $\text{AlCl}_3$ -BuPyCl mixtures. Molecular iodine does not form strong complexes in acidic or neutral solutions. Triiodide ion is stable in neutral butylpyridinium tetrachloroaluminate, but not in acidic melts. In solutions of iodine or triiodide in basic  $\text{AlCl}_3$ -BuPyCl ionic liquids equilibrium mixtures of  $\text{I}_2\text{Cl}^-$ ,  $\text{I}_3^-$ ,  $\text{ICl}_2^-$  and  $\text{I}^-$  are formed.

The charge transfer band observed for iodide solutions in neutral and basic melts reveal the formation of iodide ion - butylpyridinium cation ion pairs. Similar bands in pure basic  $\text{AlCl}_3$ -BuPyCl mixtures indicate the association of chloride ion and butylpyridinium cation.

## INTRODUCTION

Our electrochemical investigations of iodine in aluminum chloride-N-(1-butyl) pyridinium chloride (BuPyCl) ambient temperature molten salts (1,2) showed that in this solvent iodine and its compounds are involved in extensive acid-base and redox equilibria. Variations in the  $\text{AlCl}_3$ :BuPyCl mole ratio lead to wide changes in Lewis acidity of the medium. Neutral equimolar mixture of  $\text{AlCl}_3$  and BuPyCl can be made acidic by adding excess  $\text{AlCl}_3$  or basic with excess BuPyCl. The present study was undertaken to obtain additional information on interactions of iodine in different oxidation states with the ionic solvent. The electrochemical experiments did not directly identify all the iodine species formed in basic  $\text{AlCl}_3$ -BuPyCl ionic liquid (2). They did not indicate the formation of  $\text{I}_2^+$  ion in acidic ambient temperature melts although this ion has been postulated on the basis of spectral results for mixtures of  $\text{I}_2$  and  $\text{Cl}_2$  dissolved in high temperature  $\text{AlCl}_3$ -NaCl molten salts (3).

It was also of interest to seek in this medium, containing high concentration of butylpyridinium cation, charge transfer bands for iodide and chloride ions. Charge transfer, (CT), transitions for alkylpyridinium iodides have been observed in many different solvents (4-7). CT bands have also been reported for myristylpyridinium chloride solutions in chloroform (6). It was assumed (5-7) that these transitions occur for contact ion pairs; a possible contribution of solvent separated ion pairs to ionic association equilibria in these systems was however unclear (6-8). Ionic association equilibria in  $\text{AlCl}_3$ -BuPyCl ionic liquids are different in this respect in that in this medium only contact ion pairs should be formed. Formation of this type of ionic association between butylpyridinium cation and iodide and chloride ions is indicated by charge transfer bands reported herein.

#### EXPERIMENTAL

The procedures used for purification of  $\text{AlCl}_3$  (Fluka), synthesis and purification of N-(1-butyl)pyridinium chloride and preparation of the melts were similar to those described previously (9). Iodine (Malinckrodt AR), iodine chloride (Eastman Kodak), and tetraethylammonium iodide (TEAI) (Eastman Kodak) were used as received. Tetraethylammonium triiodide ( $\text{TEAI}_3$ ) was prepared according to Chattaway and Hoyle (10) and dried under vacuum.

Absorption spectra were obtained with a Perkin Elmer Model Coleman 750 spectrophotometer using 0.1 cm matched quartz cells (Wilmad). Cells were filled and sealed under a purified argon atmosphere in a Vacuum Atmosphere dry box with an HE 493 Dri Train.



## RESULTS

### Iodide and chloride ions.

Iodide ion, which in acidic melts does not exhibit any absorption in the accessible u.v. region (1), shows a single absorption band with a maxima at ca 330 nm in neutral and basic solutions (Figure 1). This band was found to obey Beer's law at iodide concentrations from  $3 \times 10^{-4}$  to  $7 \times 10^{-3}$  mol dm<sup>-3</sup> and the apparent molar absorptivity at 330 nm was  $3 \times 10^3$  dm<sup>3</sup> mol<sup>-1</sup> cm<sup>-1</sup> in the neutral as well as in basic 0.95:1 and 0.75:1 melts (Table I).

Visible or near u.v. absorption bands observed for alkylpyridinium iodides in many different solvents (4-7) have been assigned to charge transfer transitions (4-7) and taken as an evidence for contact ion pair formation (5-7). The observed independence of the apparent molar absorptivity on iodide concentration in butylpyridinium tetrachloroaluminates indicates full association of iodide ions with butylpyridinium cations. It is also interesting to note that the molar absorptivity in the ionic liquid is distinctly higher than in the other solvents where its value varied between 300 and 1600 dm<sup>3</sup> mol<sup>-1</sup> cm<sup>-1</sup> (4-7). Since in the ionic liquid only contact ion pairs are formed, the measured value can be regarded as the molar absorptivity of the contact BuPy<sup>+</sup>I<sup>-</sup> ion pair. Lower values of apparent molar absorptivity observed in the other solvents (4-7) suggest that solvent separated ion pairs are a significant fraction of ionic associations formed there.

Spectra recorded for pure melts at different AlCl<sub>3</sub>:BuPyCl mole ratios revealed a dependence of the short wavelength "cut off" on the melt composition. In acidic and neutral melts the "cut offs" were essentially

the same; an absorbance equal to 1 at 284 nm for 1.5:1 and at 285 nm for the neutral  $\text{AlCl}_3\text{:BuPyCl}$  mole ratios (1.0:1.0) melt (at 0.1 cm pathlengths, the spectra recorded versus air). For basic melt compositions these "cut offs" shifted to 313 nm for 0.95:1, 319 nm for 0.9:1 and to 325 nm for 0.75:1  $\text{AlCl}_3\text{:BuPyCl}$  mole ratios. In acidic and neutral melts the "cut offs" are due to butylpyridinium cation  $\pi\rightarrow\pi^*$  transitions (8). In aqueous solutions, maxima of alkylpyridinium cation absorption bands were observed at ca 260 nm (5). The slight difference between 1.5:1 and 1:1 melts results from an increase in  $\text{BuPy}^+$  concentration which is  $3.5 \text{ mol dm}^{-3}$  in 1.5:1 and  $4.1 \text{ mol dm}^{-3}$  in 1:1 melt. In basic melts, however, the  $\text{BuPy}^+$  concentration increase cannot account for the observed shift of the cut off;  $\text{BuPy}^+$  concentration in the 0.75:1 melt is  $4.45 \text{ mol dm}^{-3}$ . On the other hand, an absorbance edge observed between 285 and 310 nm increased when excess  $\text{BuPyCl}$  was added to the neutral butylpyridinium tetrachloroaluminate (Figure 2). The absorbance at constant wavelength varied linearly with the concentration of "free" chloride (in excess over  $\text{AlCl}_3$ ) (Figure 3). This shows that the absorption at these wavelengths is due to charge transfer transitions for a butylpyridinium cation-chloride ion pair, similar to the  $\text{BuPy}^+\dots\text{I}^-$  association. The proportionality between the absorbance and excess chloride concentration (at 305 nm Beer's law was obeyed in the chloride ion concentration range from  $5 \times 10^{-3}$  to  $0.1 \text{ mol dm}^{-3}$ ) indicates full association of butylpyridinium cation and chloride ion under the conditions employed (large excess of  $\text{RuPy}^+$ ).

The charge transfer band observed in the u.v. spectrum of  $\text{AlCl}_3\text{-RuPyCl}$  ionic liquid is a good indication of basic melt composition. It is especially useful at very low chloride concentrations, close to the neutral

butylpyridinium tetrachloroaluminate. It was found, that at concentrations below  $\sim 5 \times 10^{-3} \text{ mol dm}^{-3}$ , neither chloride nor heptachloroaluminate ions effect the electrochemical window of the molten butylpyridinium tetrachloroaluminate (2) which can be used as a criterion of the neutrality of the solvent (11). However, in u.v. spectra analogous to those shown in Figure 2,  $1 \times 10^{-3} \text{ mol dm}^{-3}$  excess chloride could be detected (at 290 nm the absorbance value corresponding to  $1 \times 10^{-3} \text{ mol dm}^{-3}$  chloride was 0.043). This method did not require the preparation of an ideally equimolar neutral melt as the reference solution. Since, as discussed above, spectra of neutral and acidic melts were essentially the same, an acidic melt could be used as the reference.

#### Iodine, iodine chloride and triiodide ion.

For solutions of iodine in neutral butylpyridinium tetrachloroaluminate a broad absorption band with a maxima at ca 480 nm ( $\epsilon_{I_2}^{480} = 800 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ ), identical to one previously observed in acidic solutions containing iodine (1), was found. A difference between the position of this band for iodine solution in liquid butylpyridinium tetrachloroaluminate and in  $\text{CCl}_4$  ( $\epsilon_{\text{max}}^{I_2} = 517 \text{ nm}$  (12)) reflects some weak intermolecular interactions between iodine molecules and solvent ions; nevertheless iodine in neutral as well as acidic  $\text{AlCl}_3\text{-BuPyCl}$  mixtures does not form strong complexes and is dissolved in the form of  $\text{I}_2$  molecules.

Spectra obtained for iodine solutions in basic  $\text{AlCl}_3\text{-BuPyCl}$  ionic liquids (Figure 4) indicated the formation of polyhalogen ions; the uncomplexed iodine band at 480 nm was not observed. The dependence of the spectra on concentration of iodine and melt composition indicated

equilibria involving chloride ion. The solvent "cut off" did not allow a direct observation of polyhalogen ion absorption maxima at wavelengths shorter than 320 nm. Only the band at ca 365 nm due to triiodide ion (vide infra) could be observed (Figure 4). Its formation in solutions of iodine dissolved in basic  $\text{AlCl}_3\text{-BuPyCl}$  melts shows that  $\text{I}_2$  is involved in some redox reaction. Reversible changes of the spectra at varying melt compositions, an independence of the spectra on time, and the fact that more than 90% of the amount of iodine initially added to a basic melt could be observed by rotating disc voltammetry after the melt was made acidic by  $\text{AlCl}_3$  addition (2), showed that the possible reactions of iodine, reduction by chloride ion and/or solvent impurities as well as the reaction with the organic cation could not account for the observed reaction. On the other hand, electrochemical experiments showed the presence of iodine (+1) as well as iodide in the solutions of iodine in basic melts (2), thus indicating that the redox reaction involving iodine is the disproportionation equilibrium



Triiodide ion is formed in the consecutive equilibrium:



Similar equilibria have been reported for  $\text{I}_2\text{Cl}^-$  solutions in acetonitrile (13), and in water (14) and recently for  $\text{I}_2/\text{CsCl-LiCl}$  melts (15). Since in the acidic melts the iodine disproportionation equilibrium is shifted far to the left (1), the addition of  $\text{AlCl}_3$  to the equilibrium mixture of polyhalogen ions existing in the solution of iodine in basic  $\text{AlCl}_3\text{-BuPyCl}$  melt causes the formation of molecular iodine.

Iodine chloride reacts with the acidic melt and is partly reduced to

iodine (1), so the spectra obtained for ICl solutions in acidic AlCl<sub>3</sub>-BuPyCl melts consisted of bands due to ICl as well as I<sub>2</sub>. Since the molar absorptivity of iodine is five times larger than the absorptivity of iodine chloride (16), and the two bands appear in the same wavelength range (the ICl band maximum was reported to be at 343 to 466 nm depending on solvent used (16)), only one very broad band was observed. The maximum of this band shifted from ca 450 to 490 nm (in 2:1 AlCl<sub>3</sub>:BuPyCl melt) over a period of 24 hours reflecting the progress of the reaction



The spectra recorded for ICl solutions in acidic AlCl<sub>3</sub>-BuPyCl mixtures were different than the spectrum reported for equimolar mixture of iodine and chlorine dissolved in the 2:1 AlCl<sub>3</sub>-NaCl melt (3). No band in the 600-800 nm region was observed; thus there was no evidence for the I<sub>2</sub><sup>+</sup> ion formation under the employed conditions. For ICl<sub>2</sub><sup>-</sup> ion two absorption bands were reported (12,13,14,15), the main one at ca. 230 nm ( $\epsilon_{\text{ICl}_2^-}^{230} = \text{ca. } 5.5 \times 10^4 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ ) and the second, much less intense band at ca 340 nm ( $\epsilon_{\text{ICl}_2^-}^{340} = \text{ca. } 300 \text{ cm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ ). The spectra obtained for

iodine chloride solutions in basic AlCl<sub>3</sub>-BuPyCl melts indicated the formation of ICl<sub>2</sub><sup>-</sup>; however only the less intense band at 338 nm could be observed. Additionally, for the latter solutions a small maximum at ca. 420 nm was evident. This band decreased at higher chloride concentrations; it probably reflected the presence of the small equilibrium concentrations of molecular ICl in the solution.

The spectra obtained for TEAl<sub>3</sub> solutions at melt compositions close to the neutral (as judged by the electrochemical window of the solvent (11)), were qualitatively similar to the spectra recorded in acetonitrile,

with two absorption maxima at 295 and 365 nm. However, apparent molar absorptivities were found to be very sensitive to small additions of  $\text{AlCl}_3$  and  $\text{BuPyCl}$  to the solution. These variations reflect changes in the conditional stability of  $\text{I}_3^-$  ion, which result from acid-base interactions of iodide with  $\text{AlCl}_3(1)$  or iodine with chloride ion (vide supra). Even at concentrations of  $\text{Cl}^-$  or  $\text{Al}_2\text{Cl}_7^-$  ions below  $5 \times 10^{-3} \text{ mol dm}^{-3}$ , which did not affect the electrochemical window of the solvent, the apparent molar absorptivity of  $\text{I}_3^-$  distinctly changed. When the melt composition was adjusted to a true "neutral" state, however, the molar absorptivity of  $\text{I}_3^-$  in neutral butylpyridinium tetrachloroaluminate ( $\epsilon_{\text{I}_3^-}^{365} = 2.25 \times 10^4 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ ) agreed well with literature data (5,13). The value of the  $\text{I}_3^-$  ion molar absorptivity in neutral butylpyridinium tetrachloroaluminate confirmed the high stability of  $\text{I}_3^-$  ion observed in electrochemical experiments (2).

Shapes of the spectra recorded for  $\text{TEAl}_3^-$  solutions in basic solvent compositions were dependent on triiodide and/or chloride ion concentrations (Figure 5). These variations were consistent with the reaction described by equilibrium 2. Thus reactions 1 and 2 lead to the formation of the equilibrium mixtures of all the ions:  $\text{I}_2\text{Cl}^-$ ,  $\text{ICl}_2^-$ ,  $\text{I}_3^-$ ,  $\text{I}^-$  and  $\text{Cl}^-$  in basic  $\text{AlCl}_3$ - $\text{BuPyCl}$  ionic liquids. In triiodide solutions reaction 2 is followed by reaction 1 (the presence of  $\text{ICl}_2^-$  and  $\text{I}^-$  ions in triiodide solutions in basic  $\text{AlCl}_3$ - $\text{BuPyCl}$  melts was observed in electrochemical experiments (2)). On the other hand, in iodine solutions iodide ion formed in reaction 1 reacts with  $\text{I}_2\text{Cl}^-$  ion according to equilibrium 2.

## DISCUSSION

U.v.-visible spectrophotometric studies characterized the species formed by iodine in different oxidation states at widely varied acidities of the  $\text{AlCl}_3\text{-BuPyCl}$  ionic liquid. The summary of these results, which are in agreement with electrochemical experiments (1,2), is shown in Table 1. The variety of species formed at different solvent acidities reflects characteristic features of the acid-base interactions in chloroaluminate molten salts. The ionic nature of the solvent promotes strong interactions between Lewis acids and chloride ion in basic melt compositions and between Lewis bases and heptachloroaluminate ion present in acidic melts. At the same time the high stability of triiodide ion in neutral butylpyridinium tetrachloroaluminate indicates relatively weak intermolecular interactions in this solvent.

Charge transfer transitions observed for butylpyridinium cation - iodide and butylpyridinium cation - chloride ion pairs indicate a high degree of association between these ions in basic  $\text{AlCl}_3\text{-BuPyCl}$  ionic liquids. The results obtained in the solutions containing a large excess of  $\text{BuPy}^+$  over  $\text{Cl}^-$  and  $\text{I}^-$  did not allow a quantitative analysis of the association equilibria. Qualitatively, however, the results observed for basic  $\text{AlCl}_3\text{-BuPyCl}$  mixtures agreed with the conclusions drawn from the analysis of the conductivity measurements for ambient temperature chloroaluminate ionic liquids (18). The proportionality between the absorbance at 305 nm and chloride ion concentration observed up to  $0.1 \text{ mol dm}^{-3}$  indicates that at formal concentrations of butylpyridinium cation up to  $4 \text{ mol dm}^{-3}$ , at least 90% of the chloride is involved in the charge transfer transition, thus is associated with  $\text{BuPy}^+$  cation (accuracy of

the measurements was arbitrarily assumed to be  $\pm 10\%$ ). This gave an estimated value for the association constant

$$K = \frac{[\text{RuPy} \cdots \text{Cl}^-]}{[\text{BuPy}^+][\text{Cl}^-]} > 2.3, \text{ which favorably compares with the value of}$$

the association constant for a similar low temperature chloroaluminate system, aluminum chloride - imidazolium chloride, which was estimated as 2 (18). The ionic association of butylpyridinium cation with chloride ion has been indicated by NMR (19,20,21) and FTIR (22) studies of  $\text{AlCl}_3\text{:BuPyCl}$  melts.

For solutions of iodine and  $\text{TEAl}_3$  in basic  $\text{AlCl}_3\text{-BuPyCl}$  ionic liquids the spectrophotometric studies confirmed polyhalogen equilibria deduced from the electrochemical results (2). In particular, the formation of  $\text{I}_3^-$  in iodine solution, expected from the values of equilibrium constant for reactions 1 and 2 estimated in the electrochemical studies (2), was directly observed. Attempts to obtain more accurate estimates of the equilibrium constants for reactions 1 and 2 from the spectral data were unfortunately unsuccessful. Spectra recorded for  $\text{I}_2$  and  $\text{TEAl}_3$  solutions in basic  $\text{AlCl}_3\text{-BuPyCl}$  mixtures were limited to a narrow range of wavelengths and consisted of overlapping bands of all ions involved in the polyhalogen equation, i.e.,  $\text{ICl}_2^-$ ,  $\text{I}_2\text{Cl}^-$ ,  $\text{I}_3^-$  and  $\text{I}^-$ . Additionally a spectrum for  $\text{I}_2\text{Cl}^-$  ion could not be obtained separately, thus molar absorptivity values for this ion were uncertain. However, the spectra obtained for iodine and triiodide solutions at low chloride concentrations (Figures 4 and 5) confirmed the approximate values of the equilibrium constants for reactions 1 and 2. Concentrations of  $\text{I}_3^-$  ions calculated from these equilibria agreed reasonably well with the  $\text{I}_3^-$  concentrations estimated from the absorbance measured of 365 nm.



-12-

CREDIT

This work was supported in part by the office of Naval Research and the Air Force Office of Scientific Research.

REFERENCES

1. Karpinski, Z. J.; Osteryoung, R. A. J. Electroanal. Chem., 1984, 164, 281.
2. Karpinski, Z. J.; Osteryoung, R. A. J. Electroanal. Chem., in press.
3. Tanemoto, K.; Mamantov, G.; Marassi, R.; Begun, G. M. J. Inorg. Nucl. Chem., 1981, 43, 1779.
4. Kosower, E. M. J. Am. Chem. Soc., 1958, 80, 3253.
5. Kosower, E. M.; Skorcz, J. Z. J. Am. Chem. Soc., 1960, 82, 2195.
6. Ray, A.; Mukerjee, P. J. Phys. Chem., 1966, 70, 2138.
7. Mackay, R. A.; Poziomek, E. J. J. Am. Chem. Soc., 1970, 92, 2432.
8. Smid, J. "Ions and Ion Pairs in Organic Reactions", p. 95, Szwarc, M. Ed., Interscience, 1972.
9. Robinson, J.; Osteryoung, R. A. J. Am. Chem. Soc., 1979, 101, 323.
10. Chattaway, F. D.; Hoyle, G. J. Chem. Soc., 1923, 123, 654.
11. Lipsztajn, M.; Osteryoung, R. A. J. Electrochem. Soc., 1983, 130, 1968.
12. Keefer, R. M.; Allen, T. L. J. Chem. Phys., 1956, 25, 1059.
13. Popov, A. I.; Swensen, R. F. J. Am. Chem. Soc., 1955, 77, 3724.
14. Cason, D. L.; Neumann, H. M. J. Am. Chem. Soc., 1961, 83, 1822.
15. Child, Jr., W. C.; Papatheodorou, G. M. J. Phys. Chem., 1983, 87, 271.
16. Bucles, R. E.; Mills, J. F. J. Am. Chem. Soc., 1954, 76, 4845.
17. Popov, A. I.; Jessup, J. N. J. Am. Chem. Soc., 1952, 74, 6127.
18. Lipsztajn, M.; Osteryoung, R. A. in preparation.
19. Tauselle, F.; Popov, A. Polyhedron, 1983, 2, 889.
20. Wilkes, J. S.; Pflug, J. L.; Levisky, J. A.; Hussey, C. L.; Fannin, Jr., A. A.; King, L. A., Frank J. Seiler Research Laboratory, FJSRL-TR-82-0005, U.S. Air Force Academy, CO.

21. Wilkes, J. S.; Levisky, J. A.; Pflug, J. L; Harvy, C. L.; Schoeffler, T. B. Anal. Chem. 1982, 54, 2379.
22. Tait, S.; Osteryoung, R. A. Inorg. Chem, in press.

Table 1

Apparent molar absorptivity of TEAI band at 330 nm in  $\text{AlCl}_3$ -RuPyCl melt.

Melt composition $N_A/N_B$	TEAI concentration $\times 10^3$ $\text{mol dm}^{-3}$	$\epsilon \times 10^{-3}$ $\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1}$
1:1	3.7	2.97
0.95:1	0.85	2.85
0.95:1	1.99	3.04
0.95:1	4.65	2.99
0.95:1	7.7	2.98
0.75	1.8	2.95
0.75	5.9	3.05

$N_A$  - Number of moles of  $\text{AlCl}_3$

$N_B$  - Number of moles of RuPyCl

TABLE 2

Iodine species at different  $\text{AlCl}_3$ -BuPyCl ionic liquid compositions.

Iodine oxidation state	Melt composition		
	Acidic	Neutral	Basic
+1	$\text{ICl}$	$\text{ICl}$	$\text{ICl}_2^-$
0	$\text{I}_2$	$\text{I}_2$	$\text{I}_2\text{Cl}^-$
-0.33	---	$\text{I}_3^-$	$\text{I}_3^-$
-1	$\text{AlCl}_3\text{I}^-$ , $\text{Al}_2\text{Cl}_6\text{I}^-$	$\text{I}^-(\text{BuPy}^+ \dots \text{I}^-)$	$\text{I}^-(\text{BuPy}^+ \dots \text{I}^-)$

FIGURE CAPTIONS

- Fig. 1. Absorption spectra of tetraethylammonium iodide solution in 0.95:1  $\text{AlCl}_3\text{:BuPyCl}$  melt.  $\text{CTEA}\text{I}$ : (1) 4.65; (2) 1.99; (3) 0.85 mM.
- Fig. 2. Absorption spectra of butylpyridinium chloride added to neutral butylpyridinium tetrachloroaluminate. Excess of  $\text{RuPyCl}$ ; (1) 18; (2) 11; (3) 5.4 mM.
- Fig. 3. Absorbances at constant wavelength measured for spectra shown in Fig. 2 plotted against chloride concentration. Wavelength: (1) 290; (2) 295; (3) 300; (4) 305 nm.
- Fig. 4. Absorption spectra of iodine in 0.95:1  $\text{AlCl}_3\text{-BuPyCl}$  melt.  $\text{CI}$ : (1) 3.5; (2) 1.68; (3) 0.85 mM.
- Fig. 5. Absorption spectra of tetraethylammonium triiodide in 0.95:1  $\text{AlCl}_3\text{-BuPyCl}$  melt.  $\text{CTEA}\text{I}_3$ : (1) 1.89; (2) 1.42; (3) 0.85; (4) 0.35mM.




FIGURE 1




FIGURE 2






FIGURE 3




FIGURE 4




FIGURE 5

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Dr. M. B. Denton  
Department of Chemistry  
University of Arizona  
Tucson, Arizona 85721

Dr. R. A. Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. J. Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. B. R. Kowalski  
Department of Chemistry  
University of Washington  
Seattle, Washington 98105

Dr. H. Freiser  
Department of Chemistry  
University of Arizona  
Tucson, Arizona 85721

Dr. H. Chernoff  
Department of Mathematics  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. A. Zirino  
Naval Undersea Center  
San Diego, California 92132

Professor George H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. Alan Bewick  
Department of Chemistry  
Southampton University  
Southampton, Hampshire  
ENGLAND 5095NA

Dr. S. P. Perone  
Lawrence Livermore Laboratory L-370  
P.O. Box 808  
Livermore, California 94550

Dr. L. Jarvis  
Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. G. M. Hieftje  
Department of Chemistry  
Indiana University  
Bloomington, Indiana 47401

Dr. Christie G. Enke  
Department of Chemistry  
Michigan State University  
East Lansing, Michigan 48824

Dr. D. L. Venezky  
Naval Research Laboratory  
Code 6130  
Washington, D.C. 20375

Walter G. Cox, Code 3632  
Naval Underwater Systems Center  
Building 148  
Newport, Rhode Island 02840

Professor Isiah M. Warner  
Department of Chemistry  
Emory University  
Atlanta, Georgia 30322

Dr. Kent Eisentraut  
Air Force Materials Laboratory  
Wright-Patterson AFB, Ohio 45433

Dr. Adolph B. Amster  
Chemistry Division  
Naval Weapons Center  
China Lake, California 93555

Dr. B. E. Douda  
Chemical Sciences Branch  
Code 50 C  
Naval Weapons Support Center  
Crane, Indiana 47322

Dr. John Eyler  
Department of Chemistry  
University of Florida  
Gainesville, Florida 32611

DL/413/83/01  
051B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Professor J. Janata  
Department of Bioengineering  
University of Utah  
Salt Lake City, Utah 84112

Dr. J. DeCorpo  
NAVSEA  
Code 05R14  
Washington, D.C. 20362

Dr. Charles Anderson  
Analytical Chemistry Division  
Athens Environmental Laboratory  
College Station Road  
Athens, Georgia 30613

Dr. Ron Flemming  
B 108 Reactor  
National Bureau of Standards  
Washington, D.C. 20234

Dr. David M. Hercules  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania

Dr. Frank Herr  
Office of Naval Research  
Code 422CB  
800 N. Quincy Street  
Arlington, Virginia 22217

Professor E. Keating  
Department of Mechanical Engineering  
U.S. Naval Academy  
Annapolis, Maryland 21401

Dr. M. H. Miller  
1133 Hampton Road  
Route 4  
U.S. Naval Academy  
Annapolis, Maryland 21401

Dr. Clifford Spiegelman  
National Bureau of Standards  
Room A337 Bldg. 101  
Washington, D.C. 20234

Dr. Denton Elliott  
AFOSR/NC  
Bolling AFB  
Washington, D.C. 20362

Dr. B. E. Spielvogel  
Inorganic and Analytical Branch  
P.O. Box 12211  
Research Triangle Park, NC 27709

Ms. Ann De Witt  
Material Science Department  
160 Fieldcrest Avenue  
Raritan Center  
Edison, New Jersey 08818

Dr. A. Harvey  
Code 6110  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. John Hoffsommer  
Naval Surface Weapons Center  
Building 30 Room 208  
Silver Spring, Maryland 20910

Mr. S. M. Hurley  
Naval Facilities Engineering Command  
Code 032P  
200 Stovall Street  
Alexandria, Virginia 22331

Ms. W. Parkhurst  
Naval Surface Weapons Center  
Code R33  
Silver Spring, Maryland 20910

Dr. M. Robertson  
Electrochemical Power Sources Division  
Code 305  
Naval Weapons Support Center  
Crane, Indiana 47522

CDR Andrew T. Zander  
10 Country Club Lane  
ONR Boston  
Plaistow, New Hampshire 03865

DL/413/83/01  
051B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Dr. Robert W. Shaw  
U.S. Army Research Office  
Box 12211  
Research Triangle Park, NC 27709

Dr. Marvin Wilkerson  
Naval Weapons Support Center  
Code 30511  
Crane, Indiana 47522

Dr. J. Wyatt  
Naval Research Laboratory  
Code 6110  
Washington, D.C. 20375

Dean William Tolles  
Naval Post Graduate School  
Spanaue1 Hall  
Monterey, California 93940

Dr. H. Wohltjen  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Robert Somoano  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California 91103

Dr. Johann A. Joebstl  
USA Mobility Equipment R&D Command  
DRDME-EC  
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus  
NASA Headquarters  
M.S. RTS-6  
Washington, D.C. 20546

Dr. Albert R. Landgrebe  
U.S. Department of Energy  
M.S. 6B025 Forrestal Building  
Washington, D.C. 20595

Dr. J. J. Brophy  
Department of Physics  
University of Utah  
Salt Lake City, Utah 84112

Dr. Charles Martin  
Department of Chemistry  
Texas A&M University  
College Station, Texas 77843

Dr. H. Tachikawa  
Department of Chemistry  
Jackson State University  
Jackson, Mississippi 39217

Dr. Theodore Beck  
Electrochemical Technology Corp.  
3935 Leary Way N.W.  
Seattle, Washington 98107

Dr. Farrell Lytle  
Boeing Engineering and  
Construction Engineers  
P.O. Box 3707  
Seattle, Washington 98124

Dr. Robert Gotscholl  
U.S. Department of Energy  
MS G-226  
Washington, D.C. 20545

Dr. Edward Fletcher  
Department of Mechanical Engineering  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. John Fontanella  
Department of Physics  
U.S. Naval Academy  
Annapolis, Maryland 21402

Dr. Martha Greenblatt  
Department of Chemistry  
Rutgers University  
New Brunswick, New Jersey 08903

Dr. John Wasson  
Syntheco, Inc.  
Rte 6 - Industrial Pike Road  
Gastonia, North Carolina 28052

Dr. Walter Roth  
Department of Physics  
State University of New York  
Albany, New York 12222

Dr. Anthony Sammells  
Eltron Research Inc.  
710 E. Ogden Avenue #108  
Naperville, Illinois 60540

Dr. W. M. Risen  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02192

Dr. C. A. Angell  
Department of Chemistry  
Purdue University  
West Lafayette, Indiana 47907

Dr. Thomas Davis  
Polymer Science and Standards  
Division  
National Bureau of Standards  
Washington, D.C. 20234



TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Paul Delahay  
Department of Chemistry  
New York University  
New York, New York 10003

Dr. P. J. Hendra  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
United Kingdom

Dr. T. Katan  
Lockheed Missiles and  
Space Co., Inc.  
P.O. Box 504  
Sunnyvale, California 94088

Dr. D. N. Bennion  
Department of Chemical Engineering  
Brigham Young University  
Provo, Utah 84602

Dr. R. A. Marcus  
Department of Chemistry  
California Institute of Technology  
Pasadena, California 91125

Mr. Joseph McCartney  
Code 7121  
Naval Ocean Systems Center  
San Diego, California 92152

Dr. J. J. Auborn  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1  
NASA-Lewis  
21000 Brookpark Road  
Cleveland, Ohio 44135

Dr. P. P. Schmidt  
Department of Chemistry  
Oakland University  
Rochester, Michigan 48063

Dr. H. Richtol  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. C. E. Mueller  
The Electrochemistry Branch  
Naval Surface Weapons Center  
White Oak Laboratory  
Silver Spring, Maryland 20910

Dr. Sam Perone  
Chemistry & Materials  
Science Department  
Lawrence Livermore National Lab.  
Livermore, California 94550

Dr. Royce W. Murray  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. G. Goodman  
Johnson Controls  
5757 North Green Bay Avenue  
Milwaukee, Wisconsin 53201

Dr. B. Brummer  
EIC Incorporated  
111 Chapel Street  
Newton, Massachusetts 02158

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Electrochimica Corporation  
Attn: Technical Library  
2485 Charleston Road  
Mountain View, California 94040

Library  
Duracell, Inc.  
Burlington, Massachusetts 01803

Dr. A. B. Ellis  
Chemistry Department  
University of Wisconsin  
Madison, Wisconsin 53706

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. M. Wrighton  
Chemistry Department  
Massachusetts Institute  
of Technology  
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons  
Department of Chemistry  
University of Utah  
Salt Lake City, Utah 84112

Donald E. Mains  
Naval Weapons Support Center  
Electrochemical Power Sources Division  
Crane, Indiana 47522

S. Ruby  
DOE (STOR)  
M.S. 6B025 Forrestal Bldg.  
Washington, D.C. 20595

Dr. A. J. Bard  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. Janet Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. Donald W. Ernst  
Naval Surface Weapons Center  
Code R-33  
White Oak Laboratory  
Silver Spring, Maryland 20910

Mr. James R. Moden  
Naval Underwater Systems Center  
Code 3632  
Newport, Rhode Island 02840

Dr. Bernard Spielvogel  
U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park, NC 27709

Dr. William Ayers  
ECD Inc.  
P.O. Box 5357  
North Branch, New Jersey 08876

Dr. M. M. Nicholson  
Electronics Research Center  
Rockwell International  
3370 Miraloma Avenue  
Anaheim, California

Dr. Michael J. Weaver  
Department of Chemistry  
Purdue University  
West Lafayette, Indiana 47907

Dr. R. David Rauh  
EIC Corporation  
111 Chapel Street  
Newton, Massachusetts 02158

Dr. Aaron Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02192

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH ENGLAND

~~Dr. R. A. Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214~~

Dr. Denton Elliott  
Air Force Office of Scientific  
Research  
Bolling AFB  
Washington, D.C. 20332

Dr. R. Nowak  
Naval Research Laboratory  
Code 6130  
Washington, D.C. 20375

Dr. D. F. Shriver  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Dr. Aaron Fletcher  
Naval Weapons Center  
Code 3852  
China Lake, California 93555

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. David Aikens  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. A. P. B. Lever  
Chemistry Department  
York University  
Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak  
Naval Ocean Systems Center  
Loc: 6343, Bayside  
San Diego, California 95152

Dr. Gregory Farrington  
Department of Materials Science  
and Engineering  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

M. L. Robertson  
Manager, Electrochemical  
and Power Sources Division  
Naval Weapons Support Center  
Crane, Indiana 47522

Dr. T. Marks  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Dr. Micha Tomkiewicz  
Department of Physics  
Brooklyn College  
Brooklyn, New York 11210

Dr. Lesser Blum  
Department of Physics  
University of Puerto Rico  
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II  
IBM Corporation  
K33/281  
5600 Cottle Road  
San Jose, California 95193

Dr. D. H. Whitmore  
Department of Materials Science  
Northwestern University  
Evanston, Illinois 60201

Dr. Alan Bewick  
Department of Chemistry  
The University of Southampton  
Southampton, SO9 5NH ENGLAND

Dr. E. Anderson  
NAVSEA-56233 NC #4  
2541 Jefferson Davis Highway  
Arlington, Virginia 20362

Dr. Bruce Dunn  
Department of Engineering &  
Applied Science  
University of California  
Los Angeles, California 90024

Dr. Elton Cairns  
Energy & Environment Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

Dr. D. Cipris  
Allied Corporation  
P.O. Box 3000R  
Morristown, New Jersey 07960

Dr. M. Philpott  
IBM Corporation  
5600 Cottle Road  
San Jose, California 95193

Dr. Donald Sandstrom  
Department of Physics  
Washington State University  
Pullman, Washington 99164

Dr. Carl Kannewurf  
Department of Electrical Engineering  
and Computer Science  
Northwestern University  
Evanston, Illinois 60201